


GETTING STARTED ON

 **LIVTENCITY**TM
(maribavir) tablets
200mg

**YOUR GUIDE TO LEARNING ABOUT
LIVTENCITY FOR THE TREATMENT
OF CMV (CYTOMEGALOVIRUS
[SAI-TOW-MEH-GUH-LOW-
VAI-RUHS])**

LIVTENCITY is a prescription medicine used to treat cytomegalovirus (CMV) infection and disease in adults and children 12 years of age and older weighing at least 77 pounds (35 kg) who have received a transplant, when their infection or disease does not respond to treatment with the medicines ganciclovir, valganciclovir, cidofovir or foscarnet. It is not known if LIVTENCITY is safe and effective in children under 12 years of age.



IMPORTANT SAFETY INFORMATION

Before taking LIVTENCITY, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if LIVTENCITY will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if LIVTENCITY passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with LIVTENCITY.

Please see additional safety information throughout, click for [Information for Patients](#), and discuss with your HCP.

WHAT IS **CMV**?



CMV is a common virus that the majority of people have been exposed to, but the immune system normally keeps in check. When the immune system cannot hold back CMV, **there are treatments** that can be used to help fight the virus.

WHY DOES CMV **MATTER TO ME**?



As someone who **received a transplant**, you may receive anti-rejection medicines to keep your immune system from attacking or rejecting the transplant. Those medicines and other medicines used before or after your transplant may leave you **vulnerable to CMV** infection and disease.



If you have an active CMV infection, it can become **refractory or resistant** to medications currently used to manage it.

- **Refractory CMV** is diagnosed when a lab test shows the level of CMV virus in the blood has not adequately decreased after 2 weeks of treatment. In addition, when a lab test shows a type of CMV with a gene mutation known to prevent the medication from working, that is considered **resistant CMV**.
- If this happens, your transplant team may then give you a different antiviral medicine in order to fight the infection.

Other considerations

Some transplant patients will experience side effects from the antiviral medicines.

- If this happens, your transplant team may decide to adjust how much of your antiviral medicine you are taking
- In some cases, your team may even change which antiviral medicines you take

WHY IS IT IMPORTANT TO **DETECT AND TREAT CMV?**




If CMV is not treated, it can lead to **negative effects on transplant outcomes**. As such, it is important to receive appropriate treatment and monitoring **with the help of your healthcare provider**.

TALK TO YOUR HEALTHCARE TEAM TO LEARN MORE

Remember, **your healthcare team is there to help you** navigate your transplant and post-transplant experience. The first step is to **talk with them**.

Here are **some topics** you may want to cover:



CMV

What are the most important things I need to know?



Your care plan

How often will I have to come in for tests or follow-up appointments?




Your feelings

How can I talk to the people in my life about what I'm going through?



Your support options

What if I need help paying for treatment?



WHAT IS **LIVTENCITY**[®] *(maribavir)?*



LIVTENCITY is a prescription medicine used to treat cytomegalovirus (CMV) infection and disease in adults and children 12 years of age and older weighing at least 77 pounds (35 kg) who have received a transplant, when their infection or disease does not respond to treatment with the medicines ganciclovir, valganciclovir, cidofovir or foscarnet. It is not known if LIVTENCITY is safe and effective in children under 12 years of age.

IS LIVTENCITY APPROVED BY THE FDA?



The **FDA approved** LIVTENCITY based on a study in adult patients who had developed a CMV infection after either a solid organ or blood stem cell transplant and had infections that were refractory with or without confirmed resistance to one or more medicines used in the study.*

*The commonly used medicines were: ganciclovir, valganciclovir, cidofovir and foscarnet.


IMPORTANT SAFETY INFORMATION (cont.)

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LIVTENCITY may affect the way other medicines work, and other medicines may affect how LIVTENCITY works and cause serious side effects.

Please see additional safety information throughout, click for [Information for Patients](#), and discuss with your HCP.

HOW SHOULD I TAKE **LIVTENCITY** (maribavir)?

2X
DAILY



TAKE 2 TIMES A DAY

WITH OR WITHOUT FOOD



SWALLOW TABLETS WHOLE



IF YOU CAN'T SWALLOW TABLETS WHOLE,

you can break apart or crush tablets, mix with drinking water, and drink.

You can also break apart tablets and take through a nasogastric (NG) or orogastric (OG) tube.

See the “Instructions for Use” in the Prescribing Information for detailed instructions.



Take LIVTENCITY exactly as your healthcare provider tells you to take it.

If you take too much LIVTENCITY, call your healthcare provider or go to the nearest hospital emergency room right away.

WHAT CAN I EXPECT WHILE **TAKING** **LIVTENCITY** (*maribavir*)?

SOME PEOPLE MAY EXPERIENCE SIDE EFFECTS WHILE TAKING LIVTENCITY.



The most common side effects while taking LIVTENCITY include changes in taste, nausea, diarrhea, vomiting, and tiredness.



Changes in taste were the most common side effect; this occurred in less than half of the patients in the trial. Changes in taste most often lasted 43 days while taking LIVTENCITY, and most often lasted 6 days after patients stopped taking LIVTENCITY.



These are not all the possible side effects of LIVTENCITY; you should always **call your doctor and transplant team for medical advice** about side effects.

NOTES

IMPORTANT SAFETY INFORMATION

What is LIVTENCITY?

LIVTENCITY is a prescription medicine used to treat cytomegalovirus (CMV) infection and disease in adults and children 12 years of age and older weighing at least 77 pounds (35 kg) who have received a transplant, when their infection or disease does not respond to treatment with the medicines ganciclovir, valganciclovir, cidofovir or foscarnet. It is not known if LIVTENCITY is safe and effective in children under 12 years of age.

IMPORTANT SAFETY INFORMATION

Before taking LIVTENCITY, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if LIVTENCITY will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if LIVTENCITY passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with LIVTENCITY.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. LIVTENCITY may affect the way other medicines work, and other medicines may affect how LIVTENCITY works and cause serious side effects.

Especially tell your healthcare provider if you take a seizure (anticonvulsant) medicine.

- **Do not start a new medicine without telling your healthcare provider.** Your healthcare provider will tell you if it is safe to take LIVTENCITY with other medicines.

What are the possible side effects of LIVTENCITY?

The most common side effects of LIVTENCITY include:

- changes in taste
- diarrhea
- tiredness
- nausea
- vomiting

These are not all the possible side effects of LIVTENCITY. Call your doctor for medical advice about side effects.

For additional safety information, click for [Patient Information](#) and discuss with your HCP.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



We're here to help.

If you've been prescribed LIVTENCITY® (maribavir), Takeda Patient Support is here to help you access your medication, find educational resources, and understand financial assistance options.

Call **1-855-268-1825** Monday-Friday, 8 AM to 8 PM ET

Our **Patient Support Managers (PSMs)** will:



Help expedite delivery of your medication



Work with your specialty pharmacy



Connect you to co-pay or insurance help



Help you understand your condition and LIVTENCITY

Call Takeda Patient Support at **1-855-268-1825**
Monday-Friday, 8 AM to 8 PM ET.

[TakedaPatientSupport.com](https://www.TakedaPatientSupport.com)



LIVTENCITY® and the LIVTENCITY logo™ are trademarks or registered trademarks of Takeda Pharmaceuticals International AG. Takeda®, the Takeda logo® and the Takeda Patient Support logo™ are trademarks or registered trademarks of Takeda Pharmaceutical Company Limited. ©2023 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 1-877-TAKEDA-7 (1-877-825-3327)
US-MAR-0486v2.0 02/23